



TITLE:

THRESHOLDS AND RESONANCES OF SCHRÖDINGER OPERATORS ON A LATTICE (Mathematical Aspects of Quantum Fields and Related Topics)

AUTHOR(S):

Hiroshima, Fumio

CITATION:

Hiroshima, Fumio. THRESHOLDS AND RESONANCES OF SCHRÖDINGER OPERATORS ON A LATTICE (Mathematical Aspects of Quantum Fields and Related Topics). 数理解析研究所講究録 2018, 2089: 146-165

ISSUE DATE:

2018-08

URL:

<http://hdl.handle.net/2433/251613>

RIGHT:

THRESHOLDS AND RESONANCES OF SCHRÖDINGER OPERATORS ON A LATTICE

Fumio Hiroshima,
Faculty of Mathematics, Kyushu University

1 Discrete Schrödinger operators $H_{\lambda\mu}$ on lattice

This is the joint work with Z. Muminov and U. Kuljanov [2] which is a continuation of [3] where a discrete Schrödinger operator with single delta potential is considered. See also related paper [1]. In the present article we consider discrete Schrödinger operators with multi-delta potentials. Let \mathbb{Z}^n be the n -dimensional lattice. The Hilbert space of ℓ^2 sequences on \mathbb{Z}^n is denoted by $\ell^2(\mathbb{Z}^n)$, and we use $\ell^2_+(\mathbb{Z}^n)$ (resp. $\ell^2_-(\mathbb{Z}^n)$) to denote its subspace of all even (resp. all odd) functions. Let $T(y)$ be the shift operator by $y \in \mathbb{Z}^n$: $(T(y)f)(x) = f(x+y)$ for $f \in \ell^2(\mathbb{Z}^n)$ and $x \in \mathbb{Z}^n$. Let $\Delta = \frac{1}{2} \sum_{\substack{x \in \mathbb{Z}^n \\ |x|=1}} (T(x) - T(0))$ be the discrete Laplacian on $\ell^2(\mathbb{Z}^n)$. Thus the discrete Schrödinger operator on $\ell^2(\mathbb{Z}^n)$ is defined by


$$\hat{H}_{\lambda\mu} = -\Delta - \hat{V},$$

where the potential \hat{V} (Figure1) depends on two parameters $\lambda, \mu \in \mathbb{R}$ and satisfies

$$(\hat{V}f)(x) = \begin{cases} \mu f(x), & \text{if } x = 0 \\ \frac{\lambda}{2} f(x), & \text{if } |x| = 1 \\ 0, & \text{if } |x| > 1 \end{cases}$$

which awards $\hat{H}_{\lambda\mu}$ to be a bounded self-adjoint operator. A notation $\mathbb{T}^n = (\mathbb{R}/2\pi\mathbb{Z})^n = (-\pi, \pi]^n$ means the n -dimensional torus equipped with its Haar measure. We set $L^2(\mathbb{T}^n) = L^2$, and let L^2_+ (resp. L^2_-) denote the subspace of all even (resp. odd) functions of the Hilbert space L^2 of L^2 -functions on \mathbb{T}^n . Let $\langle \cdot, \cdot \rangle$ mean the inner product on L^2 . Let \mathcal{F} be the standard Fourier transform $\mathcal{F} : L^2 \rightarrow \ell^2(\mathbb{Z}^n)$ defined by $\mathcal{F}f(x) = (2\pi)^{-n} \int_{\mathbb{T}^n} f(\theta) e^{-ix\theta} d\theta$. Then the inverse Fourier transform is given by $\mathcal{F}^{-1}f(\theta) = \sum_{x \in \mathbb{Z}^n} f(x) e^{ix\theta}$. The Laplacian Δ in the momentum representation is defined as $\hat{\Delta} = \mathcal{F}^{-1} \Delta \mathcal{F}$, and $\hat{\Delta}$ acts as the multiplication operator: $(\hat{\Delta} \hat{f})(p) = -E(p) \hat{f}(p)$, where $E(p)$ is given by $E(p) = \sum_{j=1}^n (1 - \cos p_j)$. Set $H_0 = -\hat{\Delta}$. The operator $H_{\lambda\mu}$ acts L^2 as

$$H_{\lambda\mu} = H_0 - V,$$

Figure 1: Potential V

and V is an integral operator of convolution type

$$(Vf)(p) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{T}^n} v(p-s)f(s)ds, \quad f \in L^2.$$

Here the kernel function is $v(p) = \frac{1}{(2\pi)^{n/2}} (\mu + \lambda \sum_{i=1}^n \cos p_i)$, and it allows the potential operator V to get the representation $V = V_{\lambda\mu}^+ + V_{\lambda}^-$, where

$$V_{\lambda\mu}^+ = \mu \langle \cdot, c_0 \rangle c_0 + \frac{\lambda}{2} \sum_{j=1}^n \langle \cdot, c_j \rangle c_j, \quad V_{\lambda}^- = \frac{\lambda}{2} \sum_{j=1}^n \langle \cdot, s_j \rangle s_j.$$

Here $\{c_0, c_j, s_j : j = 1, \dots, n\}$ is an orthonormal system in L^2 , where

$$c_0(p) = \frac{1}{(2\pi)^{n/2}}, \quad c_j(p) = \frac{\sqrt{2}}{(2\pi)^{n/2}} \cos p_j, \quad s_j(p) = \frac{\sqrt{2}}{(2\pi)^{n/2}} \sin p_j, \quad j = 1, \dots, n.$$

Adopting $V = V_{\lambda\mu}^+ + V_{\lambda}^-$, we can see that the restriction $H_{\lambda\mu}^+$ (resp. H_{λ}^-) of the operator $H_{\lambda\mu}$ to L_+^2 (resp. L_-^2) acts with the form

$$H_{\lambda\mu}^+ = H_0 - V_{\lambda\mu}^+ \quad (\text{resp. } H_{\lambda}^- = H_0 - V_{\lambda}^-).$$

Hence $H_{\lambda\mu}$ is decomposed as $H_{\lambda\mu} = H_{\lambda\mu}^+ \oplus H_{\lambda}^-$ under $L^2 = L_+^2 \oplus L_-^2$. We have $\sigma_{\text{ess}}(H_{\lambda\mu}) = \sigma_{\text{ac}}(H_{\lambda\mu}) = [0, 2n]$. Then we are interested in considering point spectrum of $H_{\lambda\mu}$ and studying their behaviors as two parameters λ and μ are varied.

2 Spectrum of even part $H_{\lambda\mu}^+$

The Birman-Schwinger principle helps us to reduce the problem to the study of spectrum of a finite dimensional linear operator. Denote by $(H_0 - z)^{-1}$ the resolvent of H_0 , where $z \in \mathbb{C} \setminus [0, 2n]$. We can see that $(H_0 - z)^{-1}V_{\lambda\mu}^+$ is a finite rank operator. Let M_{n+1} denote the linear hull of $\{c_0, \dots, c_n\}$. Then M_{n+1} is an $(n+1)$ -dimensional subspace of L_+^2 . We define $\tilde{M}_{n+1} = (H_0 - z)^{-1}M_{n+1}$ for $z \in \mathbb{C} \setminus [0, 2n]$. Then \tilde{M}_{n+1} is also an $(n+1)$ -dimensional subspace of L_+^2 since $(H_0 - z)^{-1}$ is invertible. We define $C_1 : \mathbb{C}^{n+1} \rightarrow L_+^2$ by the map

$$C_1 : \mathbb{C}^{n+1} \ni (w_0, \dots, w_n)^T \mapsto (H_0 - z)^{-1} \left(\mu w_0 c_0 + \frac{\lambda}{2} \sum_{j=1}^n w_j c_j \right) \in \tilde{M}_{n+1},$$

and define $C_2 : L_+^2 \rightarrow \mathbb{C}^{n+1}$ by

$$C_2 : L_+^2 \ni \phi \mapsto (\langle \phi, c_0 \rangle, \dots, \langle \phi, c_n \rangle)^T \in \mathbb{C}^{n+1}.$$

Then we have the sequence of maps: $L_+^2 \xrightarrow{C_2} \mathbb{C}^{n+1} \xrightarrow{C_1} L_+^2$. In particular

$$(H_0 - z)^{-1} V_{\lambda\mu}^+ = C_1 C_2.$$

Define $G_+(z) = C_2 C_1 : \mathbb{C}^{n+1} \rightarrow \mathbb{C}^{n+1}$.

Lemma 2.1 (The Birman-Schwinger principle for $z \in \mathbb{C} \setminus [0, 2n]$)

- (1) $z \in \mathbb{C} \setminus [0, 2n]$ is an eigenvalue of $H_{\lambda\mu}^+$ if and only if $1 \in \sigma(G_+(z))$.
- (2) If $z \in \mathbb{C} \setminus [0, 2n]$ and (λ, μ) satisfy $\det(G_+(z) - I) = 0$. Then $Z = (w_0, \dots, w_n)^T \in \mathbb{C}^{n+1}$ is an eigenvector of $G_+(z)$ associated with eigenvalue 1 if and only if $f = C_1 Z$, i.e.

$$f(p) = \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p) - z} \left(\mu w_0 + \frac{\lambda}{\sqrt{2}} \sum_{j=1}^n w_j \cos p_j \right) \quad (2.1)$$

is an eigenfunction of $H_{\lambda\mu}^+$ associated with eigenvalue z .

We consider the Birman-Schwinger principle for $z = 0$, which is the edge of the continuous spectrum of $H_{\lambda\mu}^+$, and it is the main issue to specify whether it is eigenvalue or threshold of $H_{\lambda\mu}^+$. In order to discuss $z = 0$ we extend the eigenvalue equation $H_{\lambda\mu}^+ f = 0$ in L_+^2 to that in L_+^1 . Note that $L_+^2 \subset L_+^1$. We consider the equation

$$E(p)f(p) - \frac{\mu}{(2\pi)^n} \int_{\mathbb{T}^n} f(p) dp - \frac{\lambda}{(2\pi)^n} \sum_{j=1}^n \cos p_j \int_{\mathbb{T}^n} \cos p_j f(p) dp = 0 \quad (2.2)$$

in the Banach space L_+^1 . Conveniently, we describe (2.2) as $H_{\lambda\mu}^+ f = 0$. Since we consider a solution $f \in L_+^1$, the integrals $\int_{\mathbb{T}^n} f(p) dp$ and $\int_{\mathbb{T}^n} \cos p_j f(p) dp$ are finite for $j = 1, \dots, n$. The unique singular point of $1/E(p)$ is $p = 0$, and in the neighborhood of $p = 0$, we have $E(p) \approx |p|^2$. Then the following is fundamental, and its proof is straightforward.

Lemma 2.2 Let $h(p) = \varphi(p)/E(p)$, where $\varphi \in C(\mathbb{T}^n)$. Then (1)-(5) follow.

- (1) It follows that $h \in L^2$ for $n \geq 5$, and $h \in L^1$ for $n \geq 3$.
- (2) Let $1 \leq n \leq 4$ and $h \in L^2$. Then $\varphi(0) = 0$.
- (3) Let $1 \leq n \leq 4$, $|\varphi(p)| < C|p|^{\alpha_n}$ for some $C > 0$ and $\alpha_n > \frac{4-n}{2}$. Then $h \in L^2$.
- (4) Let $n = 1, 2$ and $h \in L^1$. Then $\varphi(0) = 0$.
- (5) Let $n = 1, 2$, $|\varphi(p)| < C|p|^{\alpha_n}$ for some $C > 0$ and $\alpha_n > 2 - n$. Then $h \in L^1$.

Operator H_0^{-1} is not bounded in L_+^2 as well as in L_+^1 . It is however obvious by Lemma 2.2 and $V_{\lambda\mu}^+ f \in C(\mathbb{T}^n)$ that

$$L_+^2 \ni f \mapsto H_0^{-1} V_{\lambda\mu}^+ f \in L_+^2, \quad n \geq 5, \quad (2.3)$$

$$L_+^1 \ni f \mapsto H_0^{-1} V_{\lambda\mu}^+ f \in L_+^1, \quad n \geq 3. \quad (2.4)$$

Thus for $n \geq 3$ we can extend operators C_1 and C_2 . Let $n \geq 3$ and $Z = (w_0, \dots, w_n)^T$. $\bar{C}_1 : \mathbb{C}^{n+1} \rightarrow L_+^1$ is defined by

$$\bar{C}_1 Z = \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p)} \left(\mu w_0 + \frac{\lambda}{\sqrt{2}} \sum_{j=1}^n w_j \cos p_j \right)$$

and $\bar{C}_2 : L_+^1 \rightarrow \mathbb{C}^{n+1}$ by $\bar{C}_2 : L_+^1 \ni \phi \mapsto (\int_{\mathbb{T}^n} \phi(p) c_0 dp, \dots, \int_{\mathbb{T}^n} \phi(p) c_n(p) dp)^T \in \mathbb{C}^{n+1}$. Then $\bar{C}_1 \bar{C}_2 : L_+^1 \rightarrow L_+^1$. Consequently $G_+(0) = \bar{C}_2 \bar{C}_1 : \mathbb{C}^{n+1} \rightarrow \mathbb{C}^{n+1}$. Let $n \geq 3$. $\lim_{z \rightarrow 0} G_+(z) = G_+(0)$ and $\sigma(H_0^{-1} V_{\lambda\mu}^+) \setminus \{0\} = \sigma(G_+(0)) \setminus \{0\}$ follow.

Lemma 2.3 (Birman-Schwinger principle for $z = 0$) *Let $n \geq 3$. Then (1) and (2) follow.*

(1) *Equation $H_{\lambda\mu}^+ f = 0$ has a solution in L^1 if and only if $1 \in \sigma(G_+(0))$.*

(2) *Let $Z = (w_0, \dots, w_n)^T \in \mathbb{C}^{n+1}$ be the solution of $G_+(0)Z = Z$ if and only if $f = \bar{C}_1 Z$, i.e.*

$$f(p) = \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p)} \left(\mu w_0 + \frac{\lambda}{\sqrt{2}} \sum_{j=1}^n w_j \cos p_j \right) \quad (2.5)$$

is a solution of $H_{\lambda\mu}^+ f = 0$, where w_0, \dots, w_n are actually described by

$$w_0 = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{T}^n} f(p) dp, \quad w_j = \frac{\sqrt{2}}{(2\pi)^{n/2}} \int_{\mathbb{T}^n} f(p) \cos p_j dp, \quad j = 1, \dots, n. \quad (2.6)$$

By the Birman-Schwinger principle in what follows we focus on investigating the spectrum of $G_+(z)$. Since $G_+(z)$ is defined for $z \in (-\infty, 0)$ for $n = 1, 2$, and $z \in (-\infty, 0]$ for $n \geq 3$. Hence in this section we suppose that $z \in \begin{cases} (-\infty, 0) & n = 1, 2, \\ (-\infty, 0] & n \geq 3. \end{cases}$ As the function $E(p) = E(p_1, \dots, p_n)$ is invariant with respect to the permutations of its arguments p_1, \dots, p_n , the integrals used for studying the spectrum of $G_+(z)$:

$$a(z) = \langle c_0, (H_0 - z)^{-1} c_0 \rangle, \quad b(z) = \frac{1}{\sqrt{2}} \langle c_0, (H_0 - z)^{-1} c_j \rangle \quad (2.7)$$

$$c(z) = \frac{1}{2} \langle c_j, (H_0 - z)^{-1} c_j \rangle, \quad d(z) = \frac{1}{2} \langle c_i, (H_0 - z)^{-1} c_j \rangle, \quad i \neq j, \quad (2.8)$$

$$s(z) = \frac{1}{2} \langle s_j, (H_0 - z)^{-1} s_j \rangle. \quad (2.9)$$

also do not depend on the particular choice of indices $0 \leq i, j \leq n$. Note that $a(z), b(z), c(z)$ and $s(z)$ are defined for $n \geq 1$ but $d(z)$ for $n \geq 2$. Hence the $(n+1) \times (n+1)$ matrix $G_+(z)$ has the form

$$G_+(z) = \begin{pmatrix} \mu a(z) & \frac{\lambda}{\sqrt{2}} b(z) & \dots & \dots & \frac{\lambda}{\sqrt{2}} b(z) \\ \sqrt{2} \mu b(z) & \lambda c(z) & \lambda d(z) & \dots & \lambda d(z) \\ \vdots & \lambda d(z) & \ddots & \dots & \vdots \\ \vdots & \vdots & \dots & \ddots & \lambda d(z) \\ \sqrt{2} \mu b(z) & \lambda d(z) & \dots & \lambda d(z) & \lambda c(z) \end{pmatrix}. \quad (2.10)$$

In order to study the eigenvalue 1 of $G_+(z)$ we calculate the determinant of $G_+(z) - I$. We have $\det(G_+(z) - I) = \delta_r(\lambda, \mu; z) \delta_c(\lambda; z)$, where

$$\delta_r(\lambda, \mu; z) = \begin{cases} (1 - \mu a(z)) \left\{ 1 - \lambda (c(z) + (n-1)d(z)) \right\} - n \lambda \mu b^2(z), & n \geq 2 \\ (1 - \mu a(z))(1 - \lambda c(z)) - \lambda \mu b^2(z), & n = 1, \end{cases} \quad (2.11)$$

$$\delta_c(\lambda; z) = \begin{cases} \{ \lambda (c(z) - d(z)) - 1 \}^{n-1}, & n \geq 2 \\ 1, & n = 1. \end{cases} \quad (2.12)$$

We set

$$\alpha(z) = \begin{cases} c(z) + (n-1)d(z), & n \geq 2 \\ c(z), & n = 1, \end{cases} \quad \gamma(z) = a(z)\alpha(z) - nb^2(z). \quad (2.13)$$

Functions $a(z)$, $\alpha(z)$, $\gamma(z)$, $b(z)$, $c(z) - d(z)$ and $s(z)$ are monotone increasing and positive in $(-\infty, 0]$. Moreover, their limits tend to zero as z tends to $-\infty$. Note that $c(z) - d(z)$ is considered only in the case of $n \geq 2$. The following relations hold:

$$\begin{aligned} a(z)s(z) &= b(z), & n = 1, z < 0, \\ a(z)s(z) &< b(z), & n = 2, z < 0, \\ a(z)s(z) &< b(z), & n \geq 3, z \leq 0, \\ c(z) - d(z) &< s(z), & n \geq 2, z \leq 0. \end{aligned} \quad (2.14)$$

The function $a(z)/b(z)$ is monotone decreasing in $(-\infty, 0]$, and there exist limits:


$$\lim_{z \rightarrow -\infty} \frac{a(z)}{b(z)} = +\infty, \quad \lim_{z \rightarrow 0-} \frac{a(z)}{b(z)} = \begin{cases} 1, & \text{for } n = 1, 2, \\ \frac{a(0)}{b(0)}, & \text{for } n \geq 3. \end{cases} \quad (2.15)$$

We extend $\delta_r(\lambda, \mu; \cdot)$ and $\delta_c(\lambda; \cdot)$, and discuss zeros of them to specify the eigenvalue of $H_{\lambda\mu}^+$. Let $z \in (-\infty, 0)$. Applying notation in (2.13), we describe $\delta_r(\lambda, \mu; z)$ as

$$\delta_r(\lambda, \mu; z) = \gamma(z) \mathbb{H}_z(\lambda, \mu) \quad (2.16)$$

where

$$\mathbb{H}_z(\lambda, \mu) = \left(\lambda - \frac{a(z)}{b(z)} \right) \left(\mu - (n - z) \right) - n. \quad (2.17)$$

Figure 2: Hyperbola $\bar{\mathbb{H}}_z$

Instead of the equation $\delta_r(\lambda, \mu; z) = 0$, relation (2.16) allows us to study the family of rectangular hyperbola \mathbb{H}_z indexed by z , i.e. equilateral hyperbola \mathbb{H}_z on (λ, μ) -plane, which is defined by

$$\mathbb{H}_z = \{(\lambda, \mu) \in \mathbb{R}^2 \mid \mathbb{H}_z(\lambda, \mu) = 0\}$$


with asymptote $(\lambda_\infty(z), \mu_\infty(z)) = (a(z)/b(z), n - z)$. $\mathbb{H}_z(\lambda, \mu)$ can be extended to $z \in (-\infty, 0]$ for any dimension $n \geq 1$ as

$$\bar{\mathbb{H}}_z(\lambda, \mu) = \begin{cases} \mathbb{H}_z(\lambda, \mu), & z < 0, \\ (\lambda - X)(\mu - n) - n, & z = 0. \end{cases} \quad (2.18)$$

Here $X = 1$ for $n = 1, 2$ and $X = a(0)/b(0)$ for $n \geq 3$. Refer to see Figure 2. Note that $\bar{\mathbb{H}}_z(0, 0) = \frac{1}{b(z)} > 0$ for $z < 0$. We also extend the family of hyperbola \mathbb{H}_z , $z \in (-\infty, 0)$, to that of hyperbola $\bar{\mathbb{H}}_z$ indexed by $z \in (-\infty, 0]$ by

$$\bar{\mathbb{H}}_z = \{(\lambda, \mu) \in \mathbb{R} \times \mathbb{R} \mid \bar{\mathbb{H}}_z(\lambda, \mu) = 0\}.$$

For any $z_1 < z_2$, $z_1, z_2 \in (-\infty, 0]$, we note that the hyperbola $\bar{\mathbb{H}}_{z_1}$ can be moved to $\bar{\mathbb{H}}_{z_2}$ in parallel by the vector $g = \begin{pmatrix} \lambda_\infty(z_2) - \lambda_\infty(z_1) \\ \mu_\infty(z_2) - \mu_\infty(z_1) \end{pmatrix}$ whose components are positive. Refer to see Figure 3

Figure 3: Hyperbola $\bar{\mathbb{H}}_z$ moves as z approaches to $-\infty$ from 0.

Let $\beta_l(z)$ (resp. $\beta_r(z)$) denote the left brunch (resp. the right brunch) of the hyperbola $\bar{\mathbb{H}}_z$, i.e. $\bar{\mathbb{H}}_z = \beta_l(z) \cup \beta_r(z)$, where \cup denotes the disjoint union. We then see that for any $z_2 < z_1 \leq 0$ it follows that $\beta_l(z_1) \cap \beta_l(z_2) = \emptyset$ and $\beta_r(z_1) \cap \beta_r(z_2) = \emptyset$.

Lemma 2.4 *It follows that*

$$\begin{aligned} (n=1) \quad \lim_{z \rightarrow 0-} \delta_r(\lambda, \mu; z) &= \begin{cases} \infty & (\lambda, \mu) \notin \bar{\mathbb{H}}_0 \\ 1 - \mu & (\lambda, \mu) \in \bar{\mathbb{H}}_0 \end{cases}, \\ (n=2) \quad \lim_{z \rightarrow 0-} \delta_r(\lambda, \mu; z) &= \begin{cases} \infty & (\lambda, \mu) \notin \bar{\mathbb{H}}_0 \\ 1 - \mu/2 & (\lambda, \mu) \in \bar{\mathbb{H}}_0 \end{cases}, \\ (n \geq 3) \quad \lim_{z \rightarrow 0-} \delta_r(\lambda, \mu; z) &= b(0)\bar{\mathbb{H}}_0(\lambda, \mu). \end{aligned}$$

Proof: In the case of $n \geq 3$ it is trivial to see that $\lim_{z \rightarrow 0-} \delta_r(\lambda, \mu; z) = b(0)\bar{\mathbb{H}}_0(\lambda, \mu)$. Then we consider cases of $n = 1, 2$. We recall that

$$\delta_r(\lambda, \mu; z) = \gamma(z)\bar{\mathbb{H}}_z(\lambda, \mu) = \gamma(z)\bar{\mathbb{H}}_0(\lambda, \mu) + \gamma(z)(\bar{\mathbb{H}}_z(\lambda, \mu) - \bar{\mathbb{H}}_0(\lambda, \mu))$$

and

$$\gamma(z) = b(z) = a(z) - \frac{1}{n} - \frac{1}{n}za(z).$$

We can also directly see that for $n = 1, 2$

$$\bar{\mathbb{H}}_z(\lambda, \mu) - \bar{\mathbb{H}}_0(\lambda, \mu) = -\frac{1}{nb(z)}(1 + za(z))(\mu - n) + z(\lambda - \frac{a(z)}{b(z)}).$$

Together with them we have

$$\delta_r(\lambda, \mu; z) = (a(z) - \frac{1 + za(z)}{n})\bar{\mathbb{H}}_0(\lambda, \mu) + \frac{a(z)}{b(z)}(1 + za(z))(\frac{n - \mu}{n}) + \xi,$$

where

$$\xi = -za(z)(\lambda - \frac{a(z)}{b(z)}) + \frac{1 + za(z)}{n} \left(\frac{(1 + za(z))(\mu - n)}{nb(z)} + z(\lambda - \frac{a(z)}{b(z)}) \right).$$

It is well known in [5] that

$$\begin{aligned} a(z) &= \frac{1}{\sqrt{-z}\sqrt{2-z}}, \quad n = 1, \\ a(z) &= -\frac{\sqrt{2}}{2\pi} \ln(-z) + \left(\frac{1}{2} - \frac{\sqrt{2}}{\pi}\right) + O(-z), \quad \text{as } z \rightarrow 0-, \quad n = 2. \end{aligned}$$

By this it is crucial to see that

$$\lim_{z \rightarrow 0-} za(z) = 0, \quad \lim_{z \rightarrow 0-} b(z) = \infty, \quad \lim_{z \rightarrow 0-} \frac{a(z)}{b(z)} = 1$$

and

$$\lim_{z \rightarrow 0-} \xi = 0, \quad \lim_{z \rightarrow 0-} \frac{a(z)}{b(z)}(1 + za(z))(\frac{n - \mu}{n}) = 1 - \frac{\mu}{n}$$

for $n = 1, 2$. Let $n = 1$. Then

$$\delta_r(\lambda, \mu; z) = (a(z) - 1 - za(z))\bar{\mathbb{H}}_0(\lambda, \mu) + \frac{a(z)}{b(z)}(1 + za(z))(1 - \mu) + \xi$$

and the corollary follows for $n = 1$. Let $n = 2$. In a similar manner to the case of $n = 1$ we have

$$\delta_r(\lambda, \mu; z) = (a(z) - \frac{1 + za(z)}{2})\bar{\mathbb{H}}_0(\lambda, \mu) + \frac{a(z)}{b(z)}(1 + za(z))(1 - \frac{\mu}{2}) + \xi,$$

and the corollary follows for $n = 2$. Hence the proof of the corollary can be derived.

We define $\bar{\delta}_r(\lambda, \mu; z)$ for $z \in (-\infty, 0]$ by

$$\bar{\delta}_r(\lambda, \mu; z) = \begin{cases} \delta_r(\lambda, \mu; z), & z \in (-\infty, 0), \\ \lim_{z \rightarrow 0^-} \delta_r(\lambda, \mu; z), & z = 0. \end{cases} \quad (2.19)$$

From Lemma 2.4 we can see that $\bar{\delta}_r(\lambda, \mu; z)$ converges to

$$\bar{\delta}_r(\lambda, \mu; 0) = \begin{cases} 1 - \mu, & n = 1, (\lambda, \mu) \in \bar{\mathbb{H}}_0, \\ 1 - \mu/2, & n = 2, (\lambda, \mu) \in \bar{\mathbb{H}}_0, \\ 0, & n \geq 3, (\lambda, \mu) \in \bar{\mathbb{H}}_0. \end{cases} \quad (2.20)$$

Then we can show the continuity of $\bar{\delta}_r(\lambda, \mu; z)$ on z :

Lemma 2.5 *It follows that*

$(n = 1, 2)$ $\bar{\delta}_r(\lambda, \mu; z)$ is continuous in $z \in (-\infty, 0]$ for $(\lambda, \mu) \in \bar{\mathbb{H}}_0$,

$(n \geq 3)$ $\bar{\delta}_r(\lambda, \mu; z)$ is continuous in $z \in (-\infty, 0]$ for $(\lambda, \mu) \in \mathbb{R}^2$.

We set $\beta_l = \beta_l(0)$ and $\beta_r = \beta_r(0)$. The branches β_l and β_r of the hyperbola $\bar{\mathbb{H}}_0$ split \mathbb{R}^2 into three open sets

$$\begin{aligned} G_0 &= \{(\lambda, \mu) \in \mathbb{R}^2 \mid \bar{\mathbb{H}}_0(\lambda, \mu) > 0, \lambda < \lambda_\infty(0)\}, \\ G_1 &= \{(\lambda, \mu) \in \mathbb{R}^2 \mid \bar{\mathbb{H}}_0(\lambda, \mu) < 0\}, \\ G_2 &= \{(\lambda, \mu) \in \mathbb{R}^2 \mid \bar{\mathbb{H}}_0(\lambda, \mu) > 0, \lambda > \lambda_\infty(0)\}. \end{aligned}$$

Refer to see Figure 4. Hence $\partial G_0 = \beta_l$ and $\partial G_2 = \beta_r$ follow.

Lemma 2.6 (1) 1. Let $(\lambda, \mu) \in G_0 \cup \beta_l$. Then $\bar{\delta}_r(\lambda, \mu; z) \neq 0$ for $z \in (-\infty, 0)$.

2. Let $(\lambda, \mu) \in \beta_l$. Then $\bar{\delta}_r(\lambda, \mu; 0) \neq 0$ for $n = 1, 2$.

3. Let $(\lambda, \mu) \in \beta_l$. Then $\bar{\delta}_r(\lambda, \mu; 0) = 0$ for $n \geq 3$.

(2) 1. Let $(\lambda, \mu) \in G_1 \cup \beta_r$. Then $\exists_1 z \in (-\infty, 0)$ such that $\bar{\delta}_r(\lambda, \mu; z) = 0$.

2. Let $(\lambda, \mu) \in \beta_r$. Then $\bar{\delta}_r(\lambda, \mu; 0) \neq 0$ for $n = 1, 2$.

3. Let $(\lambda, \mu) \in \beta_r$. Then $\bar{\delta}_r(\lambda, \mu; 0) = 0$ for $n \geq 3$.

(3) Let $(\lambda, \mu) \in G_2$. Then $\exists_{z_1, z_2} \in (-\infty, 0)$ such that $\bar{\delta}_r(\lambda, \mu; z_1) = \bar{\delta}_r(\lambda, \mu; z_2) = 0$.

Proof: Let $(\lambda, \mu) \in G_0 \cup \beta_l$. Then we can see that $(\lambda, \mu) \notin \mathbb{H}_z$ for any $z \in (-\infty, 0)$. Thus (1)1 follows. Also it can be seen that $\cup_{z \in (-\infty, 0)} \beta_l(z) \supset G_1$, $\beta_l(z) \cap \beta_l(w) = \emptyset$ if $z \neq w$, and $\beta_r(z) \cap G_1 = \emptyset$. Hence there exists a unique $z \in (-\infty, 0)$ such that $(\lambda, \mu) \in \beta_l(z)$, which proves (2)1. We can also see that $\cup_{z \in (-\infty, 0)} \beta_l(z) \supset G_2$, $\cup_{z \in (-\infty, 0)} \beta_r(z) \supset G_2$, $\beta_l(z) \cap \beta_l(w) = \emptyset$ if $z \neq w$, $\beta_r(z) \cap \beta_r(w) = \emptyset$ if $z \neq w$, and $\beta_l(z) \cap \beta_r(z) = \emptyset$. Hence there exist $z_1, z_2 \in (-\infty, 0)$ such that $(\lambda, \mu) \in \beta_l(z_1)$ and $(\lambda, \mu) \in \beta_r(z_2)$, which proves (3). Finally we note that since $(\lambda, \mu) \in \beta_l$ implies that $1 \neq \mu$ for $n = 1$, and $2 \neq \mu$ for $n = 2$, $\bar{\delta}_r(\lambda, \mu; 0) \neq 0$ for $n = 1, 2$ follows. Hence (1) 2 and (1) 3 follow from (2.20), and (2) 2 and (2) 3 are similarly proven. **qed**




Figure 4: Region of G_j

By virtue of Lemma 2.6, $\bar{\delta}_r(\lambda, \mu; \cdot)$ has at most two zeros in $(-\infty, 0)$.

Lemma 2.7 *Let $n \geq 1$. Then (1) and (2) follow.*

- (1) *Let $\lambda \neq 0$. We assume that $z_1, z_2 \in (-\infty, 0)$ and $\delta_r(\lambda, \mu; z_k) = 0$ (if they exist). Then $1 - \mu a(z_k) \neq 0$ and $G_+(z_k)Z_k = Z_k$ has the solutions: $Z_k = (\frac{\lambda}{\sqrt{2}} \frac{nb(z_k)}{1 - \mu a(z_k)}, 1, \dots, 1)$, $k = 1, 2$, and the corresponding eigenfunctions, $H_{\lambda\mu}^+ f_k = z f_k$, are*

$$f_k(p) = \frac{\lambda}{\sqrt{2}} \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p) - z_k} \left(\mu \frac{nb(z_k)}{1 - \mu a(z_k)} + \sum_{j=1}^n \cos p_j \right), \quad k = 1, 2. \quad (2.21)$$

- (2) *Let $\lambda = 0$. We assume that $z \in (-\infty, 0)$ and $\delta_r(0, \mu; z) = 0$. Then $1 - \mu a(z) = 0$ and $G_+(z)Z = Z$ has the solution: $Z = (1, \sqrt{2}\mu b(z), \dots, \sqrt{2}\mu b(z))^T$ and the corresponding eigenfunction, $H_{\lambda\mu}^+ f = z f$, is*

$$f(p) = \frac{\mu}{(2\pi)^{n/2}} \frac{1}{E(p) - z} \quad (2.22)$$

Proof: We prove the case of $n \geq 2$. The proof for the case of $n = 1$ is similar. Since $\delta_r(\lambda, \mu; z) = 0$, we see that

$$(1 - \mu a(z)) \left(1 - \lambda (c(z) + (n-1)d(z)) \right) - n\lambda \mu b^2(z) = 0.$$

Then $1 - \mu a(z) \neq 0$ if and only if $\lambda \neq 0$, and we also have the algebraic relation

$$1 - \lambda(c(z) + (n-1)d(z)) = \frac{n\lambda\mu b^2(z)}{1 - \mu a(z)}.$$

From this relation it follows that $G_+(z_k)Z_k = Z_k$ for $\lambda \neq 0$. In the case of $\lambda = 0$ we can prove the lemma in a similar way. **qed**

We study zeros of $\delta_c(\lambda; z)$. In a similar manner we extend $\delta_c(\lambda, z)$ for $z \in (-\infty, 0]$. When $n \geq 2$, the function $c(z) - d(z)$ exists, and we can define $\alpha = \lim_{z \rightarrow 0-} c(z) - d(z)$. Note that $\alpha > 0$ and we set $\lambda_c = \frac{1}{\alpha}$. Let us write $\delta_c(\lambda; z) = \rho(\lambda; z)^{n-1}$, where $\rho(\lambda; z) = \lambda(c(z) - d(z)) - 1$. We define $\bar{\delta}_c(\lambda; z)$ by

$$\bar{\delta}_c(\lambda; z) = \begin{cases} \delta_c(\lambda; z), & z \in (-\infty, 0), n \geq 1, \\ (\lambda\alpha - 1)^{n-1}, & z = 0, n \geq 2, \\ 1, & z = 0, n = 1. \end{cases}$$

Lemma 2.8 *Let $n \geq 2$. Then (1)–(3) follow. (1) Let $\lambda \leq \lambda_c$. Then $\rho(\lambda; z) \neq 0$ for any $z \in (-\infty, 0)$. (2) Let $\lambda = \lambda_c$. Then $\rho(\lambda; 0) = 0$. (3) Let $\lambda > \lambda_c$. Then there exists unique $z \in (-\infty, 0)$ such that $\rho(\lambda; z) = 0$ with multiplicity one.*

Proof: Since $c(z) - d(z) > 0$ is strictly monotone increasing in $(-\infty, 0)$, we get

$$\begin{aligned} \rho(\lambda; z) &\leq \rho(\lambda_c; z) < \rho(\lambda_c; 0) = 0, \quad \text{if } 0 < \lambda \leq \lambda_c, \\ \rho(\lambda; z) &= -1, \quad \text{if } \lambda = 0, \end{aligned}$$

which prove (1) and (2). Since $\rho(\lambda; 0) > \rho(\lambda_c; 0) = 0$ and $\lim_{z \rightarrow -\infty} \rho(\lambda; z) = -1$ there exists $z \in (-\infty, 0)$ such that $\rho(\lambda; z) = 0$. By the monotonicity of $\rho(\lambda; \cdot)$ this zero is a unique and has multiplicity one. Hence (3) is proven. **qed**

We immediately have a lemma.

Lemma 2.9 *Let $n \geq 2$. Then (1)–(3) follow.*

- (1) *For any $\lambda \leq \lambda_c$, $\bar{\delta}_c(\lambda; \cdot)$ has no zero in $(-\infty, 0)$.*
- (2) *Let $\lambda = \lambda_c$. Then $\bar{\delta}_c(\lambda; 0) = 0$, and $z = 0$ has multiplicity $n - 1$.*
- (3) *For any $\lambda > \lambda_c$, $\bar{\delta}_c(\lambda; \cdot)$ has a unique zero in $(-\infty, 0)$ with multiplicity $n - 1$.*

Next we show the eigenfunction corresponding to zeros of $\delta_c(\lambda; \cdot)$.

Lemma 2.10 *Let $n \geq 2$, $z \in (-\infty, 0)$ and $\bar{\delta}_c(\lambda; z) = 0$. I.e., $\lambda = \frac{1}{c(z) - d(z)}$. Then the solutions*

of $G_+(z)Z = Z$ are given by $Z_j = (0, 1, 0, \dots, -1, \dots, 0)^T$, $j = 1, \dots, n - 1$, and hence the corresponding eigenfunctions, $H_{\lambda\mu}^+ g_j = z g_j$, are

$$g_j(p) = \frac{\lambda}{\sqrt{2}} \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p) - z} (\cos p_1 - \cos p_{j+1}), \quad j = 1, \dots, n - 1. \quad (2.23)$$

In particular the multiplicity of eigenvalue z is at least $n - 1$.

Now we study the spectrum located on the left edge of the essential spectrum $[0, 2n]$, i.e., $z = 0$. Suppose that $(\lambda, \mu) \in \mathbb{H}_0$. Then it is possibly $\tilde{\delta}_r(\lambda, \mu; 0) = 0$ or $\tilde{\delta}_c(\lambda, \mu; 0) = 0$. We study zeros of $\tilde{\delta}_r(\lambda, \mu; 0)$ for $n \geq 3$. We set $a(0) = a$ and $b(0) = b$, and both a and b are finite for $n \geq 3$.

Lemma 2.11 *Let $n \geq 3$. (1) Let $\lambda \neq 0$ and $\tilde{\delta}_r(\lambda, \mu; 0) = 0$. Then $1 - \mu a \neq 0$ and $G_+(0)Z = Z$ has the solution $Z = (\frac{\lambda}{\sqrt{2}} \frac{nb}{1-\mu a}, 1, \dots, 1)^T$ and the corresponding equation $H_{\lambda\mu}^+ f = 0$ has the solution:*

$$f(p) = \frac{\lambda}{\sqrt{2}} \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p)} \left(\mu \frac{nb}{1-\mu a} + \sum_{j=1}^n \cos p_j \right). \quad (2.24)$$

(2) Let $\lambda = 0$ and $\tilde{\delta}_r(0, \mu; 0) = 0$. Then $1 - \mu a = 0$ and $G_+(0)Z = Z$ has the solution: $Z = (1, 0, \dots, 0)^T$ and the corresponding equation $H_{\lambda\mu}^+ f = 0$ has the solution:

$$f(p) = \frac{\mu}{(2\pi)^{n/2}} \frac{1}{E(p)}. \quad (2.25)$$

Proof: The proof is the same as that of Lemma 2.7. **qed**

Next we show the solution corresponding to zeros of $\tilde{\delta}_c(\lambda; \cdot)$. Similar to the case of $\tilde{\delta}_r(\lambda, \mu; z) = 0$, we have the lemma below.

Lemma 2.12 *Let $n \geq 2$ and $\tilde{\delta}_c(\lambda; 0) = 0$, i.e., $\lambda = \lambda_c$. Then the solutions of $G_+(0)Z = Z$ are given by $Z_j = (0, 1, 0, \dots, \overset{j+2}{-1}, \dots, 0)^T$, $j = 1, \dots, n-1$, and hence the corresponding equation $H_{\lambda\mu}^+ g_j = 0$ has the solutions*

$$g_j(p) = \frac{\lambda_c}{\sqrt{2}} \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p)} (\cos p_1 - \cos p_{j+1}), \quad j = 1, \dots, n-1. \quad (2.26)$$

Proof: The proof is the same as that of Lemma 2.10. **qed**

As was seen above the problem for $n \geq 3$ can be reduced to study the spectrum of G_+ by the Birman-Schwinger principle, the problem for $n = 1, 2$ should be however directly investigated.

As was seen above the problem for $n \geq 3$ can be reduced to study the spectrum of G_e by the Birman-Schwinger principle, the problem for $n = 1, 2$ should be however directly investigated.

Let f be a solution of $H_{\lambda\mu}^+ f = 0$ (resp. $H_{\lambda\mu}^- f = 0$) (1) If $f \in L_+^2$ (resp. $f \in L_-^2$), we say that 0 is a threshold eigenvalue of $H_{\lambda\mu}^+$ (resp. $H_{\lambda\mu}^-$). (2) If $f \in L_+^1 \setminus L_+^2$ (resp. $f \in L_-^1 \setminus L_-^2$), we say that 0 is a threshold resonance of $H_{\lambda\mu}^+$ (resp. $H_{\lambda\mu}^-$). (3) If $f \in L_+^\varepsilon \setminus L_+^1$ (resp. $f \in L_-^\varepsilon \setminus L_-^1$) for any $0 < \varepsilon < 1$, we say that 0 is a super-threshold resonance of $H_{\lambda\mu}^+$ (resp. $H_{\lambda\mu}^-$).

Lemma 2.13 *Let $n = 1$.*

(1) Suppose that $f \in L^1(\mathbb{T})$ and $H_{\lambda\mu}^\varepsilon f = 0$. Then $f = 0$. In particular $H_{\lambda\mu}^\varepsilon$ has no threshold resonance.

- (2) There is no non-zero f such that $f \in L^\varepsilon(\mathbb{T}^2) \setminus L^1(\mathbb{T}^2)$ for some $0 < \varepsilon < 1$ and $H_{\lambda\mu}^\varepsilon f = 0$.
In particular $H_{\lambda\mu}^\varepsilon$ has no super-threshold resonance.

Proof: (1) $H_{\lambda\mu}^\varepsilon f = 0$ gives $f = \varphi/E$ and $\varphi(p) = \mu u_0 + \lambda u_1 \cos p$ by (2.2). From $f \in L^1(\mathbb{T})$ it follows that $\varphi(0) = \mu u_0 + \lambda u_1 = 0$. Hence

$$f(p) = \frac{1}{E(p)}(1 - \cos p)\mu u_0 = \mu u_0.$$

We get $u_1 = \mu u_0 \frac{1}{2\pi} \int_{\mathbb{T}} \cos t dt = 0$, which gives $\mu u_0 = 0$ and $f = 0$.

- (2) Since $f \notin L^1(\mathbb{T})$. It must be that $\mu = 0$ and $f = \varphi/E$ with $\varphi(p) = \lambda u_1 \cos p$. Hence

$$u_1 = \frac{\lambda}{(2\pi)^2} \int_{\mathbb{T}} \frac{u_1 \cos^2 p}{E(p)} dp.$$

Then $u_1 = 0$, since $\int_{\mathbb{T}} \frac{\cos^2 p}{E(p)} dp = \infty$. Then $f = 0$ follows.

Next we discuss the spectrum of $H_{\lambda\mu}^\varepsilon$ for $n = 2$ at the threshold

Lemma 2.14 Let $n = 2$.

- (1) Suppose that $f \in L^1(\mathbb{T}^2)$ and $H_{\lambda\mu}^\varepsilon f = 0$. Then $\lambda = \lambda_c$ and

$$f(p) = \lambda_c u_1 \frac{\cos p_1 - \cos p_2}{E(p)}. \quad (2.27)$$

In particular $f \in L^2(\mathbb{T}^2)$ and $H_{\lambda\mu}^\varepsilon$ has no threshold resonance.

- (2) There is no non-zero f such that $f \in L^\varepsilon(\mathbb{T}^2) \setminus L^1(\mathbb{T}^2)$ for some $0 < \varepsilon < 1$ and $H_{\lambda\mu}^\varepsilon f = 0$.
In particular $H_{\lambda\mu}^\varepsilon$ has no super-threshold resonance.

Proof: (1) Consider $H_{\lambda\mu}^\varepsilon f = 0$ in $L^1(\mathbb{T}^2)$. We can take $f = \varphi/E$ and $\varphi(p) = \mu u_0 + \lambda u_1 \cos p_1 + \lambda u_2 \cos p_2$. Since $f \in L^1(\mathbb{T}^2)$, we get $\varphi(0) = \mu u_0 + \lambda(u_1 + u_2) = 0$ and so

$$f(p) = \frac{\lambda}{E(p)}(-u_1(1 - \cos p_1) - u_2(1 - \cos p_2))$$

We obtain

$$\begin{aligned} u_1 &= -\frac{\lambda}{(2\pi)^2} \left(u_1 \int_{\mathbb{T}^2} \frac{\cos p_1(1 - \cos p_1)}{E(p)} dp + u_2 \int_{\mathbb{T}^2} \frac{\cos p_1(1 - \cos p_2)}{E(p)} dp \right), \\ u_2 &= -\frac{\lambda}{(2\pi)^2} \left(u_1 \int_{\mathbb{T}^2} \frac{\cos p_2(1 - \cos p_1)}{E(p)} dp + u_2 \int_{\mathbb{T}^2} \frac{\cos p_2(1 - \cos p_2)}{E(p)} dp \right). \end{aligned}$$

Since $\int_{\mathbb{T}^2} \frac{\cos p_1(1 - \cos p_1)}{E(p)} dp = -\int_{\mathbb{T}^2} \frac{\cos p_1(1 - \cos p_2)}{E(p)} dp$, we get

$$\begin{aligned} u_1 &= \frac{\lambda}{(2\pi)^2} (u_2 - u_1) \left(\int_{\mathbb{T}^2} \frac{\cos p_1(1 - \cos p_1)}{E(p)} dp \right), \\ u_2 &= \frac{\lambda}{(2\pi)^2} (u_1 - u_2) \left(\int_{\mathbb{T}^2} \frac{\cos p_2(1 - \cos p_2)}{E(p)} dp \right) \end{aligned}$$

and hence $u_1 = -u_2$. Consequently, $\mu u_0 = 0$, and the solution of $H_{\lambda\mu}^e f = 0$ is of the form

$$f(p) = \lambda u_1 \frac{\cos p_1 - \cos p_2}{E(p)} \in L^2(\mathbb{T}^2). \quad (2.28)$$

Inserting this into the definition of u_1 , we have $\frac{\lambda}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{\cos p_1 (\cos p_1 - \cos p_2)}{E(p)} dp = 1$ and thus taking $\lambda = \lambda_c$ we can see that (2.27) is the solution of $H_{\lambda\mu}^e f = 0$. Notice that $u_0 = 0$ follows from (2.28).

(2) Since $f \notin L^1(\mathbb{T}^2)$. It must be that $\mu = 0$ and $f = \varphi/E$ with $\varphi(p) = \lambda u_1 \cos p_1 + \lambda u_2 \cos p_2$. Hence

$$\begin{aligned} u_1 &= \frac{\lambda}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{u_1 \cos^2 p_1 + u_2 \cos p_1 \cos p_2}{E(p)} dp, \\ u_2 &= -\frac{\lambda}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{u_1 \cos p_2 \cos p_1 + u_2 \cos^2 p_2}{E(p)} dp. \end{aligned}$$

Then $u_1 = -u_2$ and $1 = \frac{\lambda}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{\cos p_1 (\cos p_1 + \cos p_2)}{E(p)} dp$. Thus $\lambda = \lambda_c$. Then f is given by (2.28), but $f \in L^2(\mathbb{T}^2)$. This contradicts with $f \notin L^1(\mathbb{T}^2)$.

Lemma 2.15 (1)-(5) follow:

- (1) Let $n = 1$. Then 0 is none of a threshold eigenvalue, a threshold resonance and a super-threshold resonance.
- (2) Let $n = 2$. Then 0 is a threshold eigenvalue with (2.26) for $(\lambda, \mu) = (\lambda_c, \mu)$ and its multiplicity is one.
- (3) Let $n = 3, 4$. Suppose $(\lambda, \mu) \in \mathbb{H}_0$. Then 0 is a threshold resonance with eigenvector (2.24) for $\lambda \neq 0$, and (2.25) for $\lambda = 0$, i.e., $(\lambda, \mu) = (0, 1/a)$.
- (4) Let $n = 3, 4$. Suppose $(\lambda, \mu) \in \mathbb{H}_0$. Then 0 is a threshold eigenvalue with (2.26) for $\lambda = \lambda_c$ and its with multiplicity is $n - 1$.
- (5) Let $n \geq 5$. Suppose $(\lambda, \mu) \in \mathbb{H}_0$. Then 0 is a threshold eigenvalue with eigenvector (2.24) for $\lambda_c \neq \lambda \neq 0$ and multiplicity one, (2.24) and (2.26) for $\lambda = \lambda_c$ and multiplicity n , and (2.25) for $\lambda = 0$, i.e., $(\lambda, \mu) = (0, 1/a)$, and multiplicity one.

Proof: (1) follows from Lemma 2.13. The solution of $H_{\lambda\mu}^+ f = 0$ is given by (2.24), (2.25) and (2.26). We note that $\int_{|p| < \varepsilon} \frac{1}{E^2(p)} dp = \infty$ for $n = 2, 3, 4$ for any $\varepsilon > 0$, and $\int_{|p| < \varepsilon} \frac{1}{E^2(p)} dp < \infty$ for $n \geq 5$ for any $\varepsilon > 0$. Since $(\lambda, \mu) \in \mathbb{H}_0$, $n \neq \mu$, and we can see that

$$\frac{nb\mu}{1-\mu a} + \sum_{j=1}^n \cos 0 = \frac{nb\mu}{1-\mu a} + n = n \left(\frac{1-\mu(a-b)}{1-\mu a} \right) = \frac{n-\mu}{1-\mu a} \neq 0.$$

Hence, using Lemma 2.2, we obtain (2.24), (2.25) $\in L^2$ for $n \geq 5$, (2.24), (2.25) $\in L^1 \setminus L^2$ for $n = 3, 4$ and (2.26) $\in L^2$ for $n \geq 2$. (2) follows from Lemmas 2.14 and 2.12. (3) and (5) follow from Lemmas 2.12 and 2.11. (4) follows from Lemma 2.12. **qed**

3 Spectrum of odd part H_λ^-

In the previous sections, we study the spectrum of $H_{\lambda\mu}^+$ by using the Birman-Schwinger principle for $n \geq 3$, and by directly solving $H_{\lambda\mu}^+ f = 0$ for $n = 1, 2$. In the case of H_λ^- we can proceed in a similar way to the case of $H_{\lambda\mu}^+$ and rather easier than that of $H_{\lambda\mu}^+$ as is seen below. Let $z \in \mathbb{C} \setminus [0, 2n]$. Let N_n be the linear hull of $\{s_1, \dots, s_n\}$. As is done for $H_{\lambda\mu}^+$, we can see that $(H_0 - z)^{-1} V_\lambda^- = S_1 S_2$. Here S_1 and S_2 are defined by

$$S_1 : \mathbb{C}^n \ni (w_1, \dots, w_n) \mapsto (H_0 - z)^{-1} \frac{\lambda}{2} \sum_{j=1}^n w_j s_j \in L_-^2,$$

$$S_2 : L_-^2 \ni \phi \mapsto (\langle \phi, s_1 \rangle, \dots, \langle \phi, s_n \rangle) \in \mathbb{C}^n.$$

We set $G_-(z) = S_2 S_1 : \mathbb{C}^n \rightarrow \mathbb{C}^n$. The following assertion is proved as Lemma 2.1.

Lemma 3.1 (Birman-Schwinger principle for $z \in \mathbb{C} \setminus [0, 2n]$)

- (1) $z \in \mathbb{C} \setminus [0, 2n]$ is an eigenvalue of H_λ^- if and only if $1 \in \sigma(G_-(z))$.
 (2) Let $z \in \mathbb{C} \setminus [0, 2n]$ and $Z = (w_1, \dots, w_n)^T \in \mathbb{C}^n$ be such that $G_-(z)Z = Z$. Then $f = S_1 Z$,

$$f(p) = \frac{1}{(2\pi)^{n/2}} \frac{1}{E(p) - z} \left(\frac{\lambda}{\sqrt{2}} \sum_{j=1}^n w_j \sin p_j \right)$$

is an eigenfunction of H_λ^- , i.e., $H_\lambda^- f = zf$.

We see that $G_-(z) = \lambda s(z)I$. Consequently we have for $n \geq 1$, $\delta_s(\lambda; z) = \det(G_-(z) - I) = (\lambda s(z) - 1)^n$. Since $G_-(z)$ is diagonal, it is very easy to find solution of $G_-(z)Z = Z$. It has n independent solutions: $Z_j = (0, \dots, \overset{j}{1}, \dots, 0)^T$. The corresponding eigenvector, $H_\lambda^- f_j = z f_j$, is given by

$$f_j(p) = \frac{1}{E(p) - z} \frac{1}{(2\pi)^{n/2}} \frac{\lambda}{\sqrt{2}} \sin p_j, \quad j = 1, \dots, n, \quad (3.1)$$

where $\lambda = 1/s(z)$. In particular the multiplicity of z is n . We can extend the Birman-Schwinger principle for $z = 0$. We extend the eigenvalue equation $H_\lambda^- f = 0$ in L_-^2 to that in L_-^1 . We consider the equation

$$E(p)f(p) - \frac{\lambda}{(2\pi)^n} \sum_{j=1}^n \sin p_j \int_{\mathbb{T}^n} \sin p_j f(p) dp = 0 \quad (3.2)$$

in L_-^1 . We also describe (3.2) as $H_\lambda^- f = 0$. We can see that $\sin p_j / E(p) \approx 1/|p|$ in the neighborhood of $p = 0$, and then $\sin p_j / E(p) \in L^1$ for $n \geq 2$. By (5) of Lemma 2.2 and $V_\lambda^- f \in C$ we can see that

$$L_-^2 \ni f \mapsto H_0^{-1} V_\lambda^+ f \in L_-^2, \quad n \geq 3, \quad (3.3)$$

$$L_-^1 \ni f \mapsto H_0^{-1} V_\lambda^- f \in L_-^1, \quad n \geq 2. \quad (3.4)$$

Thus for $n \geq 2$ we can extend operators S_1 and S_2 . Let $n \geq 2$ and $Z = (w_1, \dots, w_n)^T$. $\bar{S}_1 : \mathbb{C}^n \rightarrow L_-^1$ is defined by

$$\bar{S}_1 Z = \frac{1}{(2\pi)^{n/2}} \frac{\lambda}{\sqrt{2}} \frac{1}{E(p)} \sum_{j=1}^n w_j \sin p_j$$

and $\bar{S}_2 : L_-^1 \rightarrow \mathbb{C}^n$ by

$$\bar{S}_2 : L_-^1 \ni \phi \mapsto \left(\int_{T^n} \phi(p) s_1(p) dp, \dots, \int_{T^n} \phi(p) s_n(p) dp \right)^T \in \mathbb{C}^n.$$

Then $\bar{S}_1 \bar{S}_2 : L_-^1 \rightarrow L_-^1$. Consequently $G_-(0) = \bar{S}_2 \bar{S}_1 : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is described as an $n \times n$ matrix. Let $n \geq 2$. We have (1) $\lim_{z \rightarrow 0} G_-(z) = G_-(0)$, and (2) $\sigma(H_0^{-1} V_\lambda^-) \setminus \{0\} = \sigma(G_-(0)) \setminus \{0\}$. Hence for $n \geq 2$,

$$G_-(0) = \lambda s(0) I \quad (3.5)$$

and $\bar{\delta}_s(\lambda; z)$ is defined by

$$\bar{\delta}_s(\lambda; z) = \begin{cases} \delta_s(\lambda; z) & z \in (-\infty, 0), \\ (\lambda s(0) - 1)^n & z = 0. \end{cases} \quad (3.6)$$

Lemma 3.2 (Birman-Schwinger principle for $z = 0$) Let $n \geq 2$. Then (1) and (2) follow. (1) Equation $H_\lambda^- f = 0$ has a solution in L^1 if and only if $1 \in \sigma(G_-(0))$. (2) Let $Z = (w_1, \dots, w_n)^T \in \mathbb{C}^n$ be the solution of $G_-(0)Z = Z$ if and only if $f = \bar{S}_1 Z$, i.e.,

$$f(p) = \frac{1}{(2\pi)^n} \frac{1}{E(p)} \frac{\lambda}{\sqrt{2}} \sum_{j=1}^n w_j \sin p_j$$

is a solution of $H_\lambda^- f = 0$, where $w_j = \frac{\sqrt{2}}{(2\pi)^{n/2}} \int_{T^n} f(p) \sin p_j dp$, $j = 1, \dots, n$.

Proof: The proof is the same as that of Lemma 2.3. qed

Set $\lambda_s = \frac{1}{s(0)}$.

Lemma 3.3 Let $n \geq 1$. Then (1)-(3) follow:

- (1) Let $\lambda \leq \lambda_s$. Then $\bar{\delta}_s(\lambda; \cdot)$ has no zero in $(-\infty, 0)$.
- (2) Let $\lambda = \lambda_s$. Then $\bar{\delta}_s(\lambda_s; 0) = 0$ and $z = 0$ has multiplicity n .
- (3) Let $\lambda > \lambda_s$. Then $\bar{\delta}_s(\lambda; \cdot)$ has a unique zero in $(-\infty, 0)$ with multiplicity n .

Proof: The proof is similar to that of Lemma 2.9, and hence we omit it. qed

Threshold resonances and threshold eigenvalues for H_λ^- can be discussed by the Birman-Schwinger principle for $n \geq 2$.

Lemma 3.4 Let $n \geq 2$. Then the solutions of equation $H_\lambda^- f = 0$ are given by

$$f_j(p) = \frac{1}{(2\pi)^{n/2}} \frac{\lambda_s}{\sqrt{2}} \frac{\sin p_j}{E(p)}, \quad j = 1, \dots, n. \quad (3.7)$$

Proof: By $\bar{\delta}(\lambda_s, 0) = 0$ and Lemma 3.2 the lemma follows. **qed**

For $n = 1$ we can directly see that $H_\lambda^- f = 0$ has no solution in L^1 , but H_λ^- has a super-threshold resonance.

Proposition 3.5 (Super-threshold resonance) *Let $n = 1$. Then $H_\lambda^- f = 0$ has solution $f \in L^\varepsilon \setminus L^1$ for any $0 < \varepsilon < 1$. I.e., 0 is a super-threshold resonance of H_λ^- .*

Proof: $H_\lambda^- f = 0$ yields that

$$f(p) = \frac{\lambda}{2\pi} \frac{\sin p}{E(p)} \int_{\mathbb{T}} \sin pf(p) dp.$$

Note that however $\sin p/E(p) \notin L^1$, but we can see that $\sin p/E(p) \in L^\varepsilon$ for any $0 < \varepsilon < 1$ since $\sin p/E(p) \sim 1/p$ near $p = 0$ and $\int_{\mathbb{T}} p^{-\varepsilon} dp < \infty$. **qed**

Lemma 3.6 (1) *Let $n = 1$. Then 0 is neither a threshold resonance and a threshold eigenvalue, but for (λ, μ) with $\lambda \neq 0$, 0 is a super-threshold resonance.*

(2) *Let $n = 2$. Then 0 is a threshold resonance at $\lambda = \lambda_s$.*

(3) *Let $n \geq 3$. Then 0 is a threshold eigenvalue at $\lambda = \lambda_s$ and its multiplicity is n .*

Proof: (1) follows from Proposition 3.5. Let $n \geq 2$. Then the solution of $H_\lambda^- f = 0$ is given by (3.7). Since $\frac{\sin pj}{E(p)} \in L^1 \setminus L^2$ for $n = 2$ and $\frac{\sin pj}{E(p)} \in L^2$ for $n \geq 3$, we have $f \in L^1 \setminus L^2$ for $n = 2$, and $f \in L^2$ for $n \geq 3$. Then (2) and (3) follow. **qed**

4 Main theorems

4.1 Case of $n \geq 2$

In order to describe the main results we have to separate (λ, μ) -plane into several regions.

Lemma 4.1 *Let $n \geq 2$. Then $\lambda_\infty(z) \leq \lambda_s(z) \leq \lambda_c(z)$ for $z \in (-\infty, 0]$.*

Proof: It follows that $\lambda_c(z) = \frac{1}{c(z)-d(z)} > \lambda_s(z) = \frac{1}{s(z)} > \lambda_\infty(z) = \frac{a(z)}{b(z)}$ for $z < 0$. By a limiting argument the lemma follows. **qed**

Introduce four half planes:

$$\begin{aligned} \mathfrak{C}_- &= \{(\lambda, \mu) \in \mathbb{R}^2 | \lambda < \lambda_c\}, & \mathfrak{C}_+ &= \{(\lambda, \mu) \in \mathbb{R}^2 | \lambda > \lambda_c\} \\ \mathfrak{S}_- &= \{(\lambda, \mu) \in \mathbb{R}^2 | \lambda < \lambda_s\}, & \mathfrak{S}_+ &= \{(\lambda, \mu) \in \mathbb{R}^2 | \lambda > \lambda_s\}, \end{aligned}$$

and two vertical lines: $\beta_c = \{(\lambda, \mu) \in \mathbb{R}^2 | \lambda = \lambda_c\}$ and $\beta_s = \{(\lambda, \mu) \in \mathbb{R}^2 | \lambda = \lambda_s\}$. Note that $\mathfrak{S}_- \subset \mathfrak{C}_-$ and $\mathfrak{C}_+ \subset \mathfrak{S}_+$, and we define open sets by

$$\begin{aligned} D_0 &= G_0, & D_1 &= G_1 \cap \mathfrak{S}_-, & D_2 &= G_2 \cap \mathfrak{S}_-, & D_{n+1} &= G_1 \cap (\mathfrak{S}_+ \cap \mathfrak{C}_-), \\ D_{n+2} &= G_2 \cap (\mathfrak{S}_+ \cap \mathfrak{C}_-), & D_{2n} &= G_1 \cap \mathfrak{C}_+, & D_{2n+1} &= G_2 \cap \mathfrak{C}_+. \end{aligned}$$

	D_0	D_1	D_2	D_{n+1}	D_{n+2}	D_{2n}	D_{2n+1}
E.v.in $(-\infty, 0)$	0	1	2	$n+1$	$n+2$	$2n$	$2n+1$

	B_k	S_k	C_k	Point A	Point B
E.v. $(-\infty, 0)$	k	k	k	1	$n + 1$
0 res.	$n = 2$ —	$n = 2$ 2	$n \geq 2$ —	$n = 2$ 2	$n = 2$ —
	$n = 3, 4$ 1	$n \geq 3$ —		$n = 3, 4$ 1	$n = 3, 4$ 1
	$n \geq 5$ —			$n \geq 5$ —	$n \geq 5$ —
0.e.v.	$n = 2$ —	$n = 2$ —	$n \geq 2$ $n - 1$	$n = 2$ —	$n = 2$ 1
	$n = 3, 4$ —	$n \geq 3$ n		$n = 3, 4$ n	$n = 3, 4$ $n - 1$
	$n \geq 5$ 1			$n \geq 5$ $n + 1$	$n \geq 5$ n

Table 1: Spectrum of $H_{\lambda\mu}$ for (λ, μ) on D_k and the edges of D_k for $n \geq 2$




Figure 5: Hyperbola for $n \geq 2$

The boundaries of these sets define disjoint eight curves:

$$B_0 = \beta_l, \quad B_1 = \beta_r \cap \mathfrak{S}_-, \quad B_{n+1} = \beta_r \cap (\mathfrak{S}_+ \cap \mathfrak{C}_-), \quad B_{2n} = \beta_r \cap \mathfrak{C}_+, \\ S_1 = \beta_s \cap G_1, \quad S_2 = \beta_s \cap G_2, \quad C_{n+1} = \beta_c \cap G_1, \quad C_{n+2} = \beta_c \cap G_2,$$

and two one point sets given by $A = \beta_r \cap \beta_s$ and $B = \beta_r \cap \beta_c$.

We are now in the position to state the main theorem for $n \geq 2$.

Theorem 4.2 *Let $n \geq 2$.*

- (1) *Assume that $(\lambda, \mu) \in D_k$, $k \in \{0, 1, 2, n+1, n+2, 2n, 2n+1\}$, then $H_{\lambda\mu}$ has k eigenvalues in $(-\infty, 0)$. In addition $H_{\lambda\mu}$ has neither a threshold eigenvalue nor a threshold resonance.*
- (2) *0 is not a super-threshold resonance of $H_{\lambda\mu}$ for any $(\lambda, \mu) \in \mathbb{R}^2$.*

(3) Assume that (λ, μ) in B_k, S_k, C_k and A, B results in Table 4.1 are true:

Proof: (1) follows from Lemmas 2.6, 2.9 and 3.3. (2) follows from Lemmas 2.13 3.4 and 3.6. (3) follows from Lemmas 2.15 and 3.6. **qed**

As is drawn in Figures 5 and 9, hyperbolas for $n = 1, 2$ include the origin $(0, 0)$, which is different from those for $n \geq 3$. It actually describes that the Hamiltonian $H_{\lambda\mu}$ with $\lambda = 0$ has a negative eigenvalue for any $\mu > 0$. This is clarified in e.g., [3] and is also hold for one or two dimensional Schrödinger operator of the form $-(1/2)\Delta + V$. We refer to see e.g., [6] and reference therein. Moreover for any $(\lambda, \mu) \in (0, \infty) \times (0, \infty)$, $H_{\lambda\mu}$ always has negative eigenvalue and dose not have threshold resonance nor threshold eigenvalue for $n = 1, 2$.

By a Rellich type theorem for the discrete Schrödinger operator, embedded eigenvalues may be absent in the interval $(0, 2n)$. Under some assumptions this is proven in [4]. We are however interested in constructing a Hamiltonian $H_{\lambda\mu}$ which has two eigenvalues on both edges of the spectrum $[0, 2n]$. See Figure 6 below:



Figure 6: Two embedded eigenvalues or resonances of $H_{\lambda\mu}$

By the discussion stated in the previous sections we can construct an example of discrete Schrödinger operators which has two embedded eigenvalues. Let $n \geq 3$. Points (λ, μ) on the blue curve in the upper left construct Schrödinger operators $H_{\lambda\mu}$ which has resonance or eigenvalue at 0. On the other hand points (λ, μ) on the red curve in the upper right construct Schrödinger operators $H_{\lambda\mu}$ which has resonance or eigenvalue at $2n$. See Figure 7.




Figure 7: Resonance and eigenvalue on edges

We have the theorem.

Theorem 4.3 (1) Let $n = 3, 4$. Then there exists two pints (λ_1, μ_1) and (λ_2, μ_2) in (λ, μ) -plane such that both 0 and $2n$ of the spectrum $\sigma(H_{\lambda_j\mu_j})$ are simultaneously resonances. (2) Let $n \geq 5$. Then there exists two pints (λ_1, μ_1) and (λ_2, μ_2) in (λ, μ) -plane such that both 0 and $2n$ of the spectrum $\sigma(H_{\lambda_j\mu_j})$ are simultaneously eigenvalues.

Proof: Two branches $\beta_r(2n)$ and β_l cross at just two points (λ_1, μ_1) and (λ_2, μ_2) for $n \geq 3$, where $\lambda_1, \mu_2 < 0$ and $\lambda_2, \mu_1 > 0$. Refer to see Figure 8. In the case of $n = 3, 4$, two points

0 and $2n$ are resonances and in the case of $n \geq 5$ these are eigenvalues. Then the proof is completed. **qed**




Figure 8: Crossing points of hyperbolas

4.2 Case of $n = 1$

Let $n = 1$. In this case, the asymptote of \tilde{H}_0 is $(\lambda_\infty(0), \mu_\infty(0)) = (1, 1)$, and λ_c is not defined. We also see that $\lambda_s = 1 = \lambda_\infty(0)$. Then we have 4 sets $D_0 = G_0, D_1 = G_1 \cap \mathfrak{S}_-, D_2 = G_1 \cap A_1$ and $D_3 = G_2$. The boundaries of these sets define disjoint three curves: $B_0 = \beta_l, B_2 = \beta_r$ and $A_1 = \beta_s$.

	D_0	D_1	D_2	D_3		B_k	A_1
E.v.in $(-\infty, 0)$	0	1	2	3	E.v.in $(-\infty, 0)$	k	1
					0 res.	—	—
					0.e.v.	—	—

Table 2: Spectrum of $H_{\lambda\mu}$ for (λ, μ) on D_k and the edges of D_k for $n = 1$

Now we formulate next result for $n = 1$.

Theorem 4.4 *Let $n = 1$.*

- (1) *Assume $(\lambda, \mu) \in D_k, k \in \{0, 1, 2, 3\}$. Then $H_{\lambda\mu}$ has k eigenvalues in $(\infty, 0)$. In addition 0 is neither a threshold resonance and a threshold eigenvalue.*
- (2) *Assume that (λ, μ) with $\lambda = \lambda_s = 1$. Then 0 is a super-threshold resonance.*
- (3) *Assume that (λ, μ) in B_k or A_1 . Then results in Table 4.2 are true: In particular $H_{\lambda\mu}$ has neither a threshold resonance nor a threshold eigenvalue.*

Proof: The theorem follows from Lemmas 2.6, 2.9, 3.3, 2.15 and 3.6. **qed**

Acknowledgements This work is financially supported by Grant-in-Aid for Science Research (B)16H03942 from JSPS. We thank kind hospitality of Euler International Mathematical Institute, St. Petersburg, Russia, and E.Korotyaev for organizing "2nd Summer School: Various aspects of mathematical physics" held in July 8-11, 2017, where this work is partial done. We also thank J. Møller, Y. Nomura and H. Morioka to send us attention of Theorem 4.3. Finally we thank Tomoko Eto for drawing beautiful Figures 1-8.

